

Effectiveness of Foot Reflexology on Blood Pressure, Headache and Fatigue Among Hypertensive Patients at Tertiary Care Hospital, Karad

Corresponding Author:

Mr. Ajay Jyotiram Kawar

Assistant Professor

Dept. Of Medical-Surgical Nursing

Hirai Institute Of Nursing Education Malwadi, Masur.

ABSTRACT: Background:- As per the WHO statistics (2019) an estimated 1.13 billion people worldwide have hypertension, most (two-thirds) living in low and middle income countries. In 2015, 1 in 4 men and 1 in 5 women had hypertension. Fewer than 1 in 5 people with hypertension have the problem under control. Hypertension is a major cause of premature death worldwide. One of the global targets for non-communicable diseases is to reduce the prevalence of hypertension by 25% by 2025.

PROBLEM STATEMENT EFFECTIVENESS OF FOOT REFLEXOLOGY ON BLOOD PRESSURE, HEADACHE AND FATIGUE AMONG HYPERTENSIVE PATIENTS AT TERTIARY CARE HOSPITAL, KARAD

OBJECTIVES: -To find out association between sociodemographic variables with blood pressure

- To find out association between sociodemographic variables with headache

-To find out association between sociodemographic variables with fatigue

METHODS: A quantitative approach was used to conduct among 60 patients which were selected by simple random sampling technique. 30 samples were allotted in Experimental group and 30 in control group at Medicine ward and ICU in Krishna Hospital, karad. The experimental group received Foot Reflexology Once a day for Five Days and each session lasted for 20 minutes where control group followed hospital routine management. The data was analyzed by using descriptive and inferential statistics.

RESULTS: Pretest the mean and SD of SBP was 164 ± 16.30 in the experimental group and 164.87 ± 16.97 in the control group. Pretest the mean and SD of DBP was 100.73 ± 6.46 in the experimental group and 101.53 ± 7.68 in the control group. Among the Experimental group in the pretest, the Mean and SD of SBP was 164 ± 16.30 and DBP was 100.73 ± 6.46 . In posttest, the Mean and SD of SBP was 155.47 ± 15.96 and that of 101.53 ± 7.68 . The test of Significance was calculated using paired "t" test. The calculated "t" value for SBP was $2.116 (P=0.0430)$. And that of DBP was $2.305 (p=0.285)$ were greater than the table value. It showed that Foot Reflexology was effective in reducing the high blood pressure among the patients with hypertension. The Mean post-test value of SBP was 155.47 ± 15.96 in the experimental group and 154.2 ± 10.35 in the control group. The Mean post-test value of DBP was 96.33 ± 8.26 in the experimental group and 92.67 ± 5.10 in the control group. The calculated paired "t" value for SBP is 2.305 in the experimental group. And DBP was 5.972 in the control group. These values were greater than the table value. It showed that Foot Reflexology was effective in controlling the Blood Pressure among the patients with Hypertension. Among the experimental group, in pre-test, the Mean and SD of Headache was 4.67 ± 1.97 and in the post-test the mean and SD of Headache was 2.27 ± 2.13 . The test of significance was calculated using paired "t" test. The calculated "t" value for Headache was $4.079 (P=0.003)$ were greater than table value. It showed that Foot Reflexology was effective in reducing Headache among hypertensive patients.

Among the experimental group, in the pre-test, the Mean and SD of fatigue was 28.77 ± 5.41 and in post-test the Mean and SD of Fatigue was 27.7 ± 6.79 . The test of significance was calculated using paired "t" test. The calculated "t" value is $4.013 (p=0.004)$. It showed that Foot Reflexology was effective in reducing the level of fatigue among Hypertensive patients.

CONCLUSION: Reflexology helps overall circulation in the body and it helps to reduce the blood pressure. Hypertension is one of the conditions purported to be improved by complementary therapies such as Foot Reflexology. The investigator conducted study to determine the effectiveness of Foot Reflexology on Blood pressure, Headache and Fatigue in Hypertensive patients in tertiary care hospital, karad.

KEY WORDS: Foot Reflexology, Systolic Blood Pressure, Diastolic Blood Pressure, Headache. Fatigue.

I.INTRODUCTION:

Maintenance of good health is the means to living, existence, zest for life, feelings of being and happiness. Health not only means absence of sickness but presence of feelings and behaviors which constitutes different kinds of health. Achieving and maintaining health is an ongoing process, shaped by both the evolution of health care knowledge and practices as well as personal strategies and organized interventions for staying healthy known as lifestyle management. As per the WHO statistics (2019) an estimated 1.13 billion people worldwide have hypertension, most (two-thirds) living in low and middle income countries. In 2015, 1 in 4 men and 1 in 5 women had hypertension. Fewer than 1 in 5 people with hypertension have the problem under control. Hypertension is a major cause of premature death worldwide. One of the global targets for non-communicable diseases is to reduce the prevalence of hypertension by 25% by 2025. Hypertension is called a "silent killer". Most people with hypertension are unaware of the problem because it may have no warning signs or symptoms. For this reason, it is essential that blood pressure is measured regularly. Reducing hypertension prevents heart attack, stroke, and kidney damage, as well as other health problems.

The age-adjusted prevalence of hypertension in India was 11.3% (95% CI 11.16% to 11.43%) among persons aged between 15 and 19 and was four percentage points higher among males 13.8% ((95% CI 13.46% to 14.19%) than among females 10.9% (95% CI 10.79% to 11.06%)

Headache disorder is a major public health issue and is a great burden for the person, the health care system, and society. Headache has troubled mankind from the dawn of civilization. For millennia, the medical and popular literature has described headache triggers, relieving factors, and the signs and symptoms of the migraine complex, including headache, aura, prodromal, nausea or vomiting, and familial tendency. Headache is a common human experience, diverse in its expression, complex in its manifestation, and difficult to understand by any simple mechanism. Migraine is predominantly an affliction of young people. The strong familial association and early onset of the disorder suggest that there is an important genetic component.

Fatigue can refer to a subjective symptom of malaise and aversion to activity or to objectively impaired performance. It has both physical and mental aspects. The symptom of fatigue is a poorly defined feeling, and careful inquiry is needed to clarify complaints of "fatigue," "tiredness," or "exhaustion" and to distinguish lack of energy from loss of motivation or sleepiness, which may be pointers to specific diagnoses. The physiological and psychological mechanisms underlying subjective fatigue are poorly understood. Fatigue may rather be usefully regarded as a final common pathway for a variety of causal factors.

Hypertension is less often reported associated with fatigue although there is some evidence that hypertensive subjects suffer more fatigue-related symptoms.

Reflexology is potentially a very valuable therapeutic nursing skill and could have wide-ranging and cost-effective benefits in health care, from special care baby units through to care of the elderly. Like many other complementary therapies, reflexology seems to restore and maintain health by rebalancing the body. Whilst too many, reflexology may appear a gentle therapy, it is vital that the contraindications are known and that it is only carried out by trained therapists. The International Institute of Reflexology defines reflexology as a science that deals with the principle that there are reflex areas in the feet, and stimulating them properly can help many health problems in a natural way a type of preventative maintenance (International Institute of Reflexology, 2012). The foundations of reflexology can be traced to two different theories or Schools of thought documented in the reflexology literature. The first theory originated in traditional Chinese medicine (TCM) and the second one in a Western technique known as Zone therapy.

Reflexology is an active and continually evolving therapeutic practice. In recent years traditional reflexology has branched out into new and exciting therapeutic expressions in an effort to improve the health and wellness of clients. Within all of these different types of reflexology, there are therapists who specialize in different protocols.

NEED FOR STUDY: Globally, an estimated 26% of the world's population (972 million people) has hypertension, and the prevalence is expected to increase to 29% by 2025, driven largely by increases in economically developing nations. The high prevalence of hypertension exacts a tremendous public health burden. As a primary contributor to heart disease and stroke, the first and third leading causes of death worldwide, respectively, high blood pressure was the top modifiable risk factor for disability adjusted life-years lost worldwide in 2013.

Headache affects people of all ages, races and socioeconomic status and is more common in women. Some headaches are extremely debilitating and have significant impact on an individual's quality of life, imposing huge costs to healthcare and indirectly to the economy in general. Only a small proportion of headache disorders require specialist input. The vast majority can be effectively treated by a primary care physician or generalist with correct clinical diagnosis that requires no special investigation. Headaches are broadly divided into primary and secondary. Secondary headaches are uncommon but their recognition is extremely important as timely intervention may be lifesaving. A reflexology treatment aims to restore the body's natural balance. In this busy world, we spend a lot of time on our feet and our feet take a daily pounding. We put our feet under a lot of pressure and never give them a thought until they start hurting. This impact is felt throughout the whole body and can even lead to tension headaches.

Foot reflexology is a non-invasive, cost effective method used for the reduction of blood pressure. It is a readily available, painless procedure that can be applied to any person without consideration time and place. This form of treatment demands no special devices or requirements.

There are a number of therapies which have a positive and holistic approach in a natural way. Complementary therapy is proved to be one of the effective treatments for most of the disease conditions. Complementary therapy such as Yoga, acupuncture and homeopathy improve circulation, help boost immune system, eliminate toxins, reduce stress and tension, relieve pain, induce deep relaxation and restore balance to body systems.

PROBLEM STATEMENT "EFFECTIVENESS OF FOOT REFLEXOLOGY ON BLOOD PRESSURE, HEADACHE AND FATIGUE AMONG HYPERTENSIVE PATIENTS AT TERTIARY CARE HOSPITAL, KARAD"

OBJECTIVES:

1. To find out association between sociodemographic variables with blood Pressure.
2. To find out association between sociodemographic variables with headache.
3. To find out association between sociodemographic variables with fatigue.

HYPOTHESIS:

H₀ –There will be no significant effect of Foot reflexology On Blood Pressure, Headache and Fatigue. H₁-There will be significant effect of Foot Reflexology on Blood Pressure, Headache, Fatigue.

OPERATIONAL DEFINITIONS:

Effectiveness: It refers to the reduction of high blood pressure, headache and fatigue among hypertensive patients as measured by using sphygmomanometer, as determined by the difference in pre and post test score and which is statistically significant.

Foot Reflexology: It refers to the therapeutic application of five steps of massage to both feet of patients with hypertension for a period of 10 Minutes on each foot, daily for 5 consecutive days as a relaxations therapy.

Blood Pressure: Blood pressure is a measure of the force that your heart uses to pump blood around your body.

Headache: Headache is pain in any region of the head. Headaches may occur on one or both sides of the headbe isolated to a certain location, radiate across the head from one point, or have a viselike quality.

Fatigue: In this study, it refers to extreme tiredness resulting from mental and physical illness. **Hypertension:** In this Study It refers to blood pressure is above 140/90mmHg. **Patients with Hypertension:** In this study, it refers to the patients, diagnosed as hypertension for a period of less than 10 years.

Tertiary Care Hospital: In this study, a tertiary care hospital refers to Krishna Hospital, karad.

II. REVIEW OF LITERATURE:

1. Lee YM. (2011) The study was to evaluate the effects of self-foot reflexology on stress (perceived stress, urine cortisol level, and serum cortisol level), fatigue, skin temperature and immune response in female undergraduate students. The research design was a nonequivalent control group pretest-posttest design. Participants were 60 university students: 30 in the experiment group and 30 in the control group. The period of this study was from April to June 2010. The program was performed for 1 hr. a session, three times a week for 6 weeks. The data were analyzed using the SPSS/WIN 17.0 program. The results showed that self-foot reflexology was effective in reducing perceived stress and fatigue, and raised skin temperature in female undergraduate students. But cortisol levels and immune response were not statistically significant different. The results of this study indicate that self-foot reflexology is an effective nursing intervention in reducing perceived stress and fatigue and, in improving skin temperature. Therefore, it is recommended that this be used in clinical practice as an effective nursing intervention for in female undergraduate students.

2. Song MR, Song HM. Th (2005) this study was to examine the effects of foot reflexology on ADL and fatigue in stroke patients. Method: The subjects were 31 stroke patients hospitalized in the Oriental Medicine Hospital of D University from June to November, 2002. Foot reflexology was applied to the experimental group twice a week for 6 weeks, 40 minutes each. For the data analysis, x²-test was conducted to verify the homogeneity of general characteristics and clinical characteristics, and t-test was done to verify the homogeneity of ADL and fatigue. To examine the relative efficacy of the intervention, ANOVA and ANCOVA were conducted. Results: After foot reflexology, the subjects in the experimental group showed significant improvement in ADL. They also had less physical, psychological, and neurosensory fatigue, which are three areas of fatigue. Conclusion: The results suggest that foot reflexology is an effective intervention that helps the body work efficiently, eases stress and strain, and enhances the homeostasis of the body through stimulating the reflex zone of internal organs in the body. Therefore, it is necessary to develop foot reflexology as a unique nursing intervention.

III. RESEARCH METHODOLOGY:

Research Methodology is a technique for proficiently dealing with the research problem. It is a study of concentrating how research is done scientifically. This chapter deals with the methodology which includes research approach, research design, setting of the study, Population, the sample and the sampling techniques, development of tool, procedure of data collection and plan of data analysis.

Research Approach: A research approach mentions to the nurse researcher what information to gather and how to analyse it. It additionally proposes potential ends to be drawn from the data. In the present study, the nurse investigator evaluates the effectiveness of foot reflexology on blood pressure, headache and fatigue among hypertensive patients. It centralizing on the nature of the research problem for this study and the objective to be fulfilled, a quantitative approach was used to conduct this study.

Research Design: The research design is the plan, structure and strategy of investigations of answering the research question is the overall plan or blue print the researcher select to carry out their study. The assurance of research design depends on the purpose of the study; investigate approach and variable to be considered. The research design close for the investigation was True experimental research design.

Two groups were studied; Experimental group (Intervention) Control Group (Routine Management) Variable are qualities, Properties or characteristics of person, thing or situation that change or vary.

Independent Variable: Foot reflexology b) Dependent Variable: Blood Pressure, Headache and Fatigue.

Setting of the study: Setting is the more explicit places where data collection occurs dependent on the idea of the research question and the kind of data expected to address it. The study was conducted in Medicine ICU and Medicine Wards at Krishna Hospital, Karad.

Population: In this study, the population consisted of Hypertensive Patients those are suffering from Headache and Fatigue in the tertiary hospital, karad Sample Size The sample comprised of 60 patients with hypertension, comprising of 30 samples in experimental group and 30 control groups.

Based on this study done by, T Sasi Priya .Effectiveness of Foot Reflexology on Blood pressure among patients with hospital, Coimbatore. SBP: 4.67 ± 3.33 DBP: 9.07 ± 2.60 By SBP 1. $n = 4 / (\times)$ Sampling Technique: Polit and Hungler, 1999 defined sampling technique is the process of selecting a portion of the population to represent the entire population. Non-probability purposive sampling was selected for the present study. Sampling Technique: Polit and Hungler, 1999 defined sampling technique is the process of selecting a portion of the population to represent the entire population. Non-probability purposive sampling was selected for the present study.

Sampling Technique: Polit and Hungler, 1999 defined sampling technique is the process of selecting a portion of the population to represent the entire population. Non-probability purposive sampling was selected for the present study. Sampling Technique: Polit and Hungler, 1999 defined sampling technique is the process of selecting a portion of the population to represent the entire population. Non-probability purposive sampling was selected for the present study.

The demographic data was collected through structured interview schedule in Marathi. Blood Pressure was measured and recorded and Headache and fatigue was assessed for 5 consecutive days. Foot reflexology was implemented. Post-test blood pressure was taken for all the samples after 30 min duration of the pre-test. Ethical aspects were considered throughout the study.

IV. RESULT AND SUMMARY:

In experimental group, 16(53.33%) sample belongs to the age group of 41 to 50 years, 16(53.33%) sample were Female, most of the sample 29(96.67%) were married, 17(5.67%) samples had secondary education , 10(33.33%) sample were self-employed and doing private job, 9(30%)samples were earning monthly income of 10000-150000Rs, 19(69.33) samples were belongs to nuclear family , 12(86.67%) sample are taking mixed type of diet, 19(63.33%) samples were from Urban region, 15(60%) sample were used Tabaco products, 19(63.33) sample had hypertension for period of less than 5 years, 20(66.67%) sample had no family history of hypertension, 21(70%) samples were taken regular treatment of hypertension, 16(53.33%) samples suffered from respiratory illnesses, 23(76.67%) samples took medication of hypertension less than 5 years.

In control group , 17(56.67%)samples were in the age group of 41 to 50 years, 15(50%) samples were male and 15(50%) samples were female, 28(93.33%) samples were married, 12(40%) samples had higher secondary education, 11(36.67) samples were self-employed , half of the samples 15(50%) were monthly income of 10 to 15 thousands, 18(60%) samples belonged to nuclear family , 26(86.67) samples took mixed type of diet, 21(70) samples from rural region, 18(60%) samples used tobacco and its products, 16(60%) samples had hypertension for a period of less than 5 years, 18(60%) samples had no family history of hypertension, 23(76.67) samples took regular medication of hypertension, 17(56.67%) samples suffered from respiratory illnesses, 20(66.67%) samples took medication for less than five years. Jasvirkaur, Sukhpal Kaur, NeerjaBhardwaj (2011) conducted a study to assess the effect of foot massage and reflexology on physiological parameters of critically ill patients in Chandigarh. 60 patients admitted in various ICUs of Nehru hospital were selected for the study. Similar to the present study, this study was consistent with the demographic variables. According to age, the mean age (years) \pm SD of the subjects was 46.7 ± 16.1 , with the range of 16-80 years .Around none third (31.6%) were between 31-45 years. Majority (70%) were male. 30% were illiterate. 28.3% were self-employed. Around half (53.3%) of the subjects had 1-5 family members in their families.

To find out association between sociodemographic variables with blood Pressure.

In the experimental group with regard to SBP there was not significant association found between Age, Gender, Education, Occupation, Income per month, Marital Status, Type of Family, Diet pattern, Residence, Habits, Duration of hypertension, Family history of hypertension, Treatment of hypertension, Associated Illness and Duration of Medication. In the control group with regard to SBP there was a significant association found in the Duration of medication only and

not significant association between Age, Gender, Education, Occupation, Income per month, Marital Status, Type of Family, 99 Diet pattern, Residence, Habits, Duration of hypertension, Family history of hypertension, Treatment of hypertension and Associated Illness. In the experimental group with regard to DBP there was not significant association found between Age, Gender, Education, Occupation, Income per month, Marital Status, Type of Family, Diet pattern, Residence, Habits, Duration of hypertension, Family history of hypertension, Treatment of hypertension, Associated Illness and Duration of Medication. In the control group with regard to DBP there was significant association found in Residence only and was not significant association found between Age, Gender, Education, Occupation, Income per month, Marital Status, Type of Family, Diet pattern, Residence, Habits, Duration of hypertension, Family history of hypertension, Treatment of hypertension, Associated Illness and Duration of Medication.

To find out association between sociodemographic variables with headache.

In the experimental group with regard to Headache there was significant association found in Type of family and not significant association found between Age, Gender, Education, Occupation, Income per month, Marital Status, Diet pattern, Residence, Habits, Duration of hypertension, Family history of hypertension, Treatment of hypertension, Associated Illness and Duration of Medication. In the control group with regard to Headache there was significant association found in the Gender and not significant association found between Age, Education, Occupation, Income per month, Marital Status, Type of Family, Diet pattern, Residence, Habits, Duration of hypertension, Family history of hypertension, Treatment of hypertension, Associated Illness and Duration of Medication.

To find out association between sociodemographic variables with fatigue.

In the experimental group with regard to fatigue there was significant association found in Marital status and not significant association found between Age, Gender, Education, Occupation, Income per month, Type of Family, Diet pattern, Residence, Habits, Duration of hypertension, Family history of hypertension, Treatment of hypertension, Associated Illness and Duration of Medication. In the control group with regard to fatigue there was significant association found in the Gender and Occupation and not significant association found between Age, Education, Income per month, Marital Status, Type of Family, Diet pattern, Residence, Habits, Duration of hypertension, Family history of hypertension, Treatment of hypertension, Associated Illness and Duration of Medication.

V. SUMMARY, CONCLUSION, IMPLICATIONS AND RECOMMENDATIONS:

Summary of the study: Hypertension is defined as an average systolic blood pressure above 140 mmHg and a diastolic blood pressure above 90 mmHg or both. The first line of treatment for hypertension includes dietary changes, physical exercise, and weight loss. Reflexology helps overall circulation in the body and it helps to reduce the blood pressure. Hypertension is one of the conditions purported to be improved by complementary therapies such as Foot Reflexology. The investigator conducted study to determine the effectiveness of Foot Reflexology on Blood pressure, Headache and Fatigue in Hypertensive patients in tertiary care hospital, karad.

The Objectives of the study Was: To find out association between sociodemographic variables with fatigue.

The Following hypotheses were tested.

H0 –There will be no significant effect of Foot reflexology On Blood Pressure, Headache and Fatigue.

H1-There will be significant effect of Foot Reflexology on Blood Pressure, Headache, Fatigue.

In experimental group, 16(53.33%) sample belongs to the age group of 41 to 50 years, 16(53.33%) sample were Female, most of the sample 29(96.67%) were married, 17(5.67%) samples had secondary education , 10(33.33%) sample were self-employed and doing private job, 9(30%)samples were earning monthly income of 10000-150000Rs, 19(69.33) samples were belongs to nuclear family , 12(86.67%) sample are taking mixed type of diet, 19(63.33%) samples were from Urban region, 15(60%) sample were used Tabaco products, 19(63.33) sample had hypertension for period of less than 5 years, 20(66.67%) sample had no family history of hypertension, 21(70%) samples were taken regular treatment of hypertension, 16(53.33%) samples suffered from respiratory illnesses, 23(76.67%) samples took medication of hypertension less than 5 years.

In control group, 17(56.67%) samples were in the age group of 41 to 50 years, 15(50%) samples were male and 15(50%) samples were female, 28(93.33%) samples were married, 12(40%) samples had higher secondary education, 11(36.67%) samples were self-employed, half of the samples 15(50%) were monthly income of 10 to 15 thousands, 18(60%) samples belonged to nuclear 104 family, 26(86.67%) samples took mixed type of diet, 21(70%) samples from rural region, 18(60%) samples used tobacco and its products, 16(60%) samples had hypertension for a period of less than 5 years, 18(60%) samples had no family history of hypertension, 23(76.67%) samples took regular medication of hypertension, 17(56.67%) samples suffered from respiratory illnesses, 20(66.67%) samples took medication for less than five years.

CONCLUSION: The study was done to evaluate the effectiveness of Foot Reflexology on Blood Pressure, Headache and fatigue among hypertensive patients in tertiary care hospital, Karad. The Statistical analysis of the study showed that there was decrease in the elevated blood pressure level, Headache and fatigue level after implementation of Foot Reflexology in patients with hypertension when compared with the pretest. Thus this study proved the effectiveness of Foot Reflexology on the Blood Pressure, Headache and fatigue among patients with hypertension.

IMPLICATIONS: The findings of the study have implications in different aspects of nursing profession such as nursing practice, nursing education, nursing research and nursing administration.

Nursing Practice: • Nurses play a vital role in prevention of non-communicable diseases (NCD). The incidence and prevalence of hypertension and its complications are increasing every year. Thus, there is an urgent need to concentrate on the measures to reduce the disease burden. • Foot Reflexology can be incorporated in the daily nursing routine as it is a proven technique to reduce the elevated Blood pressure, headache and level of Fatigue. • The nursing personnel should be responsible to create awareness in the general public through mass media campaign regarding the importance of foot reflexology as an adjuvant therapy for hypertension and prevent its complications.

Nursing Education: • As a Nurse Educators, we must strengthen the non-pharmacological methods of managing hypertension and should be incorporated in nursing subjects. • Nursing education should emphasize on preparing nurses to various treatment modalities and update their knowledge in all fields including complementary and alternative medicine. • This study will enhance the nursing students to acquire knowledge about Foot Reflexology and its importance in maintaining the Blood pressure, No Headache and no fatigue. • Student nurses can be trained in participating foot Reflexology so that they can inculcate it in nursing care activities. **Nursing Research:** • This study can be a baseline for future studies to build upon and motivate the investigators to conduct further studies. • There is a need for extensive research in hypertension and its non pharmacological measures such as reiki, laughter therapy, yoga and other relaxation techniques. • As Nursing profession focuses on evidence based practice, the nursing personnel should involve in research activities to come out with successful remedies to reduce the burden of various diseases.

Nursing Administration: • Nurse administrators should organize various staff development programs to educate the nurses on importance of foot reflexology as an adjunct to manage hypertension. • Nurse administrators should motivate the nurses to gain knowledge regarding various alternative therapies for hypertension and implement them while caring the clients

RECOMMENDATIONS: The study recommends the following the further research The study can be conducted with large samples to generalize the findings. Comparative studies can be conducted between various alternative modalities like comparison of foot reflexology with reiki. The study can be conducted in different clinical settings. Comparative study can be undertaken between the genders. The same study can be conducted in community settings where the family members can be taught foot reflexology. Similar study can be conducted with longer duration of intervention.

VI. REFERENCES:

1. Silambarasan R. A Study to evaluate the effectiveness of foot relexology on blood pressure among patients with hypertention at selected hospital, Erode (Doctoral dissertation, Anbu College of Nursing, Komarapalayam).
2. World Hypertension Day 2019 [Internet]. World Health Organization. World Health Organization; 2019 [cited 2021Jun5]. Available from: https://www.who.int/cardiovascular_diseases/world-hypertension-day-2019/en/
3. A leading multidisciplinary medical journal [Internet]. BMJ Open. [cited 2021Jun5]. Available from: <https://bmjopen.bmj.com/>
4. Home - PMC - NCBI [Internet]. National Center for Biotechnology Information. U.S. National Library of Medicine; [cited 2021Jun5]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles>
5. Hindustan Times, Mumbai | By SadaguruPandit UPDATED ON MAY 17, 2019 08:45 AM IST
6. ChobanianAV;BakrisGL;BlackHR;CushmanWC;GreenLA;IzzoJL;JonesDW;MartersonBJ;OparilsS;WrightJT;Roccella EJ; ; Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure [Internet]. Hypertension (Dallas, Tex. : 1979). U.S. National Library of Medicine; [cited 2021Jun5]. <https://pubmed.ncbi.nlm.nih.gov/14656957/> Available from:
7. Peng KP, Wang SJ. Epidemiology of headache disorders in the Asia-Pacific region. *Headache: The Journal of Head and Face Pain*. 2014 Apr;54(4):610-8
8. Silberstein SD, Upton RB, Goadsby PJ. *Headache in clinical practice*. Routledge; 2018 May 8.
9. Silberstein SD, Rosenberg J. Multispecialty consensus on diagnosis and treatment of headache. *Neurology*. 2000 Apr 25;54(8):1553
10. Patient-Centered Assessment of Symptoms and Outcomes - Full Text View [Internet]. Full Text View - ClinicalTrials.gov. [cited 2021Jun5]. Available from: <https://clinicaltrials.gov/ct2/show/NCT03952624>
11. Harbison JA, Walsh S, Kenny RA. Hypertension and daytime hypotension found on ambulatory blood pressure is associated with fatigue following stroke and TIA. *QJM: An International Journal of Medicine*. 2009 Feb 1;102(2):109-15.
12. Silambarasan R. A Study to evaluate the effectiveness of foot relexology on blood pressure among patients with hypertention at selected hospital, Erode (Doctoral dissertation, Anbu College of Nursing, Komarapalayam).
13. GUNNARSDOTTIR TJ. Reflexology. specialist at the University of Minnesota Medical Center, Fairview. She is an adjunct clinical professor at the University of Minnesota School of Nursing and adjunct assistant professor at the University of Minnesota School of Medicine. Dr. Tracy has been the principal investigator or coinvestigator on. 2014:459. 113
14. *Med. J. Cairo Univ.*, Vol. 79, No. 2, September: 193-202, 2011 www.medicaljournalofcairouniversity.com
15. Reflexology Association of Canada. 2021 [cited 2021Jun5]. Available from: <https://www.reflexologycanada.org/en/>
16. What is the global prevalence of hypertension (high blood pressure)? [Internet]. Latest Medical News, Clinical Trials, Guidelines - Today on Medscape. 2020 [cited 2021Jun5]. Available <https://www.medscape.com/answers/241381-7614/what-is-the-global-prevalence-of-hypertension-high-blood-pressure> from:
17. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Dennison C, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the prevention, detection, evaluation, and management of high blood pressure in adultsexternal icon . *Hypertension*. 2018;71(19):e13–115.
18. Originally published29 2019<https://doi.org/10.1161/HYPERTENSIONAHA.119.13437> Jul *Hypertension*. 2019;74:450–457
19. Ahmed F. Headache disorders: differentiating and managing the common subtypes [Internet]. *British journal of pain*. SAGE Publications; 2012 [cited 2021Jun5]. Available https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4590146/table/table1_2049463712459691/ 114 from:
20. Suffering from a headache? This foot reflexology massage could help [Internet]. *Healthista*. 2019 [cited 2021Jun5]. Available from: <https://www.healthista.com/suffering-from-a-headache-this-foot-reflexology-massage-could-help/>
21. Lee J, Han M, Chung Y, Kim J, Choi J. Effects of foot reflexology on fatigue, sleep and pain: a systematic review and meta-analysis. *Journal of Korean Academy of Nursing*. 2011 Dec;41(6):821-33.
22. Brent A. Bauer MD. Reflexology for stress relief [Internet]. Mayo Clinic. Mayo Foundation for Medical Education and Research; 2020 [cited 2021Jun5]. Available from: <https://www.mayoclinic.org/healthy-lifestyle/consumer-health/expert-answers/what-is-reflexology/faq-20058139>

23. Footnotes A step- by-steo guide to the practice of foot reflexology by Sr. Mary EM McGlone.
24. Kotruchin P, Imoun S, Mitsungnern T, Aountrai P, Domthaisong M, Kario K. The effects of foot reflexology on blood pressure and heart rate: A randomized clinical trial in stage-2 hypertensive patients. *The Journal of Clinical Hypertension*. 2021 Mar;23(3):680-6.
25. Retnaningsih D. The Effect of Reflexology on the Blood Pressure Reduction.
26. Arslan G, Ceyhan Ö, Mollaoğlu M. The influence of foot and back massage on blood pressure and sleep quality in females with essential hypertension: a randomized controlled study. *Journal of Human Hypertension*. 2020 Jul 16:1 1. 115
27. Missiriya S, Hassan JS, Anuncia A. Effect of Reflex Therapy on Stress and Blood Pressure among Older Adults with Hypertension. *Journal of Pharmaceutical Research International*. 2020 Aug 5:110-7.
28. Elvira M. Influence of yoga on the blood pressure of elderly with hypertension. *Malaysian Journal of Medical Research (MJMR)*. 2018 Jul 2;2(3):30-5.
29. El-Abd AA, Abdalla KF, Mohamed HG, Ali MM. Effect of foot reflexology on blood pressure and quality of life among patients with essential hypertension. Published master thesis. Medical surgical nursing, faculty of nursing, Benhauniversity. 2018'
30. Eshghizade M, Mojtavavi SJ, Najafi S, Nodar HZ, Khani N. The effect of foot reflexology on blood pressure in elderly patients with hypertension. *Avicenna Journal of Phytomedicine*. 2015 Oct 2;5
31. T Priya S. Effectiveness of foot reflexology on blood pressure among patients with hypertension at selected hospital, Coimbatore (Doctoral dissertation, Kongunadu College of Nursing, Coimbatore).
32. Sujatha T, Judie A. Effectiveness of a 12-week yoga program on physiopsychological parameters in patients with hypertension. *International Journal of Pharmaceutical and Clinical Research*. 2014;6(4):329-35.
33. George R, D'silva F, D'souza JL. Perceived Barriers and Effectiveness of Planned Teaching Programme on Life Style Modification Practices of Persons 116 with Hypertension-A Study in Dakshina Kannada, Mangalore. *Journal of Krishna Institute of Medical Sciences (JKIMSU)*. 2012 Jul 1;1(2).
34. Smith TA, Thurgood SL. A pragmatic case series of clients living with medically diagnosed migraines self-referred to reflexology. *Complementary Therapies in Clinical Practice*. 2020 Nov 1;41:101230.
35. Imani N, Shams SA, Radfar M, Ghavami H, Khalkhali HR. Effect of applying reflexology massage on nitroglycerin-induced migraine-type headache: A placebo-controlled clinical trial. *Agri: Agri (Algoloji) Derneği'nin Yayın Organidir= The Journal of the Turkish Society of Algology*. 2018 Jul 1;30(3):116-22.
36. Bertrand A, Mauger-Vauglin CE, Martin S, Goy F, Delafosse C, Marec-Berard P. Evaluation of efficacy and feasibility of foot reflexology in children experiencing chronic or persistent pain. *Bulletin du cancer*. 2019 Dec 1;106(12):1073-9.
37. Eom MR, Kim SK, Park MH, Namkim O. The Effects of Foot Reflexology on Sleep Quality, Pain and Muscle Strength in Community Dwelling Female Elderly. *Indian Journal of Public Health Research & Development*. 2019 May 1;10(5).
38. Rambod M, Pasyar N, Shamsadini M. The effect of foot reflexology on fatigue, pain, and sleep quality in lymphoma patients: a clinical trial. *European Journal of Oncology Nursing*. 2019 Dec 1;43:101678.
39. Ko YS, Park MK. Effects of self-foot reflexology on fatigue and sleep states in women nurses. *Korean journal of women health nursing*. 2007;13(1):21-7.
40. Roshanravan M, Jouybari L, BahraniTaghanaki H, Vakili M, Sanagoo A, Amini Z. Effect of foot reflexology on fatigue in patients undergoing hemodialysis: a sham-controlled randomized trial. *Journal of Mazandaran University of Medical Sciences*. 2016 Jun 15;26(137):32-41.
41. Jin SJ, Kim YK. The effects of foot reflexology massage on sleep and fatigue of elderly women. *Korean Journal of Adult Nursing*. 2005;17(3):493-502.
42. Lee YM. Effects of self-foot reflexology on stress, fatigue, skin temperature and immune response in female undergraduate students. *Journal of Korean Academy of Nursing*. 2011 Feb 1;41(1):110-8.
43. Bagheri-Nesami M, Zargar N, Gholipour-Baradari A, Khalilian A. The effects of foot reflexology massage on pain and fatigue of patients after coronary artery bypass graft. *Journal of Mazandaran University of Medical Sciences*. 2012 Sep 10;22(92):52-62.
44. Song MR, Song HM. The effects of foot reflexology on ADL and fatigue in stroke patients. *The Korean Journal of Rehabilitation Nursing*. 2005;8(2):139 48.

45. Izadpanah A, Sahbaei F, Shojaei F, Fesharaki M. Effect of Foot Reflexology on Fatigue in Women Admitted to Rama Spa Massage Service Center. *Journal of Health and Care*. 2012 Nov 10;14(4):0-.
46. Bahrami T, Rejeh N, Heravi-Karimooi M, Vaismoradi M, Tadrissi SD, Sieloff CL. Aromatherapy massage versus reflexology on female elderly with acute coronary syndrome. *Nursing in critical care*. 2018 Sep;23(5):229-36.

NEWSPAPER ARTICLE:

1. Foot Reflexology: Happy Feet Therapy. *Times of India Health*
2. Prevalence of hypertension high among lower, middle class population in India. *The Times of India*.

INTERNET RESOURCES:

1. <http://pubmed.com>
2. <http://www.reflexologychartsforhighbloodpressure.com> .
3. <http://www.who.orghttp://www.bodywithsoul.com>
4. <http://www.heartfoundation.org.au> .
5. <http://www.researchgate.com> .
6. <http://www.pubmed.com> .
7. <http://www.isojournal.org> .
8. <http://www.en.wikipedia.org> .
9. <http://www.biomedcentral.com> .
10. <http://www.soothes.co.uk> .